

March 1998

Erratum: Jet Cluster Algorithms in e^+e^- Annihilation

S. Bethke

*III. Physikalisches Institut,
RWTH Aachen, Germany*

Z. Kunszt

*Institute of Theoretical Physics,
ETH Zürich, Switzerland*

D.E. Soper

*Institute of Theoretical Science, University of Oregon,
Eugene, Oregon, USA*

W.J. Stirling

*Department of Physics, University of Durham,
Durham, England*

Abstract

We correct an important misprint in the journal version of our earlier work on "*New Jet Cluster Algorithms: Next-to-leading Order QCD ...*", published in Nucl. Phys. B 370 (1992) 310, which may have lead to an incorrect parametrisation of the leading order QCD coefficients for the JADE type jet cluster algorithms.

Only recently we were made aware of a subtle misprint in the journal version of our work on "*New Jet Cluster Algorithms: Next-to-leading Order QCD and Hadronization Corrections*" [1]; the preprint version of that publication [2] is correct and does **not** contain this typing error.

The misprint in [1] appears in the first line of numbers presented in Table 2, where we gave a simple parametrisation of the QCD coefficients for jet production rates in hadronic final states of e^+e^- annihilations. In particular, the parameter k_1 parametrisng the leading order coefficient $A(y)$ for the relative production rate of 3-jet events using the JADE-type jet algorithms [3, 4],

$$f_3(y) = A(y) \frac{\alpha_s}{2\pi} ,$$

should correctly be -6.513 (as given in [2]) instead of -6.153 (as misprinted in [1]).

This misprint, if implemented in experimental determinations of α_s from measurements of jet production rates using the JADE jet finding algorithm, may result in an overestimate of α_s by 5 to 10 %, depending on the range of jet resolution parameters and centre of mass energy utilised in the analysis.

The **correct** part of Table 2 should therefore read

	Algorithm	y range	k_0	k_1	k_2	k_3	k_4
A	E, E0, p	0.01 - 0.33	3.096	-6.513	3.463	-0.157	0.0134

where $A(y)$ is parametrised according to equation 22 of [1, 2]:

$$A(y) = \sum_{n=0}^{n=4} k_n \left(\log \frac{1}{y} \right)^n .$$

The coefficients k_n for the leading- and the next-to-leading order QCD predictions of the other jet schemes given in [1, 2] are correct and are not repeated here.

Acknowledgement

We thank A. Brandenburg for making us aware of the misprint in Table 2 of [1].

References

- [1] S. Bethke, Z. Kunszt, D.E. Soper and W.J. Stirling, Nucl. Phys. B370 (1992), 310.
- [2] S. Bethke, Z. Kunszt, D.E. Soper and W.J. Stirling, CERN-TH.6222/91.
- [3] JADE collaboration, S. Bethke et al., Phys. Lett. B213 (1988) 235.
- [4] OPAL collaboration, M.Z. Akrawy et al., Z. Phys. C49 (1991) 375.